

From XRISM to NewAthena

Aurora Simionescu



XRISM proposal preparation workshop SRON, 13.02.2024


NewAthena is an ESA L-class mission that will serve as the major X-Ray observatory, world-wide, in the late 2030's and beyond.

Like XRISM, NewAthena carries an X-ray micro-calorimeter (the XIFU), and a wide field CCD imager (the WFI), but with significantly enhanced sensitivity and spatial resolution compared to Resolve and Xtend.


Athena, originally selected in 2014, underwent a successful re-design in 2022/2023, aiming to achieve compliance with the ESA long-term budget with minimal change on the science case.

NewAthena retains nearly all of the Athena science, and can also answer new questions that have emerged over the last decade. The new systems design was deemed realistic, technically and financially, and was unanimously endorsed by the SPC in November 2023.






Spectroscopy: from Resolve to XIFU





XRISM/Resolve (GVC):

- 170 cm2 effective area at 7 keV
- 1.3' PSF
- 35 pixels
- 3x3' FOV
- ~5 eV spectral resolution
- 2 12 keV bandpass

4 spatial resolution elements per FOV







NewAthena/XIFU:

- 870 cm2 effective area at 7 keV
- 9" PSF
- over 1500 calorimeter pixels
- 4' diameter FOV
- <4 eV spectral resolution
- 0.3 12 keV bandpass

>2200 spatial resolution elements per FOV

Imaging: from Xtend to WFI

XRISM/Xtend NewAthena/WFI


~440 cm2 effective area @1.5 keV

- 10000 cm2 effective area @ 1 keV


1.3' PSF - 9" PSF, stable across FOV

38x38' FOV - 40x40' FOV





From XRISM to NewAthena: the science



Galaxy and cluster evolution

Diffuse gas in local environments, Physics and dynamics of cosmic plasmas

Formation and evolution of compact objects

The lives and deaths of stars

10 Mpc

Properties of hot gas in L* galaxies

Properties of gas filling cosmic web filaments

Redshift evolution⁵

Dynamics of hot ICM


Fundamental science question

Quantify the dissipation of gravitational energy into bulk motions and turbulence of the intracluster medium


Experiment

Spatially-resolved measurement of turbulent line broadening and shifts out to R500

XRISM will start addressing this (e.g. Coma Center 200ks, A2029: 250ks at R2500!) but NewAthena will move us towards representative samples and detailed mapping.



100ks pointing centred around R500 of a fiducial z=0.15 cluster similar to RX J1720.1+2638.



True (top) and reconstructed (bottom) maps for a fiducial Coma-like simulated cluster observed with the X-IFU (Roncarelli et al. 2018).


Fundamental science question


Quantify how AGN feedback is disspated into bulk motions and turbulence

Experiment

Spatially-resolved measurement of turbulent line broadening and shifts on scales < AGN cavities

XRISM will map dynamics around larger (older) AGN lobes in a few nearby systems.





100 ks NewAthena/WFI images of 6 simulated L* galaxies showing bubble feedback in TNG50

Fundamental science question

Determine how (AGN) feedback impacts typical L* galaxies

Experiment

Image substructures in the circumgalactic medium of z~0.01 L* galaxies


Fundamental science question

Detect the extremely tenuous gas *typical* of cosmic web filaments


Experiment

Absorption spectroscopy of the Warm-Hot Intergalactic Medium using quasars as backlight

+Mapping filaments connecting to nearby clusters with WFI!



25 WHIM detections with 4 Ms [assumes 2% uncertainty on relative effective area calibration]
Credit: F. Nicastro (OAR)




Fundamental science question

Quantify chemical enrichment in the ICM since the early epochs of structure formation


Experiment

Measure abundances of both alpha- and Fe-group elements up to z=2

XRISM: excellent measurements of Si, S, Ar, Ca, Fe, Ni in local clusters; O, Ne, Mg inaccessible if GVC; high-z targets too faint even for GVO



3-sigma detection of Mg in z=2 galaxy cluster in 140 ks of NewAthena XIFU.



Black hole spin distribution

Geometry of X-ray emitting corona

Wind launching mechanisms and energetics

Growth of early SMBH at high-z


Multimessenger science and GW

Fundamental science question


Determine the nature of the primary X-ray source in Active Galactic Nuclei (AGN) and stellar compact objects

Experiment

High-resolution spectraltiming, accretion disk reverberation mapping (time lags)



Statistical constraints with 100 ks (4x25) NewAthena/X-IFU



Black hole spin vs. ionised outflow density

Black hole mass vs. corona height

Credit: P. Boorman, G. Mastroserio (CalTech), O. König (FAU)


Fundamental science question

Determine how powerful AGN outflows are launched and how they impact the evolution of galaxies

Experiment

Track evolution of disk outflows on their dynamical time-scale; probe launch mechanism via line shapes

XRISM requires integrating signal over longer exposure times.




Fundamental science question

Determine the space density of AGN that dominate black hole growth up to the EoR

Experiment

Measure the population of moderately obscured (NH=1e22 cm-2) AGN up to z~8



Conservative predictions for 31.6 Ms survey covering 40 deg2. Recent JWST results show space density may be much higher than assumed.


Credit: J. Aird (UoE), F. Carrera (IFCA)

Fundamental science question

Multi-messenger astrophysics (e.g., counterparts of neutron star merging gravitational wave sources up to z~1)

Experiment

Light curves can be followed for a longer time and at greater angles from the jet allowing more EM counterpart detections




The lives and deaths of stars

Stellar flare dynamics and (exo)planet habitability

Supernova progenitors, explosion mechanisms, and particle acceleration in SNR

Neutron star equation of state




The lives and deaths of stars


Fundamental science question

Constrain the equation of state of neutron stars

Experiment

Measure with %-level accuracy the radius of rotationallypowered pulsars through pulse profile modelling





Credit: S. Guillot (IRAP), N. Rea (CSIC), T. Salmi, S. Vinciguerra, A. Watts (UvA)

The lives and deaths of stars (and planets!)


Fundamental science question

Determine if and how stellar coronal activity (may) impact planet atmospheres

Experiment

Dynamics of stellar coronae, estimate of coronal mass losses. Compare flares between planet hosting and non-planet hosting stars.

Spectral evolution of stellar flares in their own dynamical timescales (~500 s) NewAthena/XIFU simulation of young (40 Myr) planet hosting star DS Tuc A Credit: R. Osten (STScI)



The lives and deaths of stars

Fundamental science questions

Constrain SN explosion mechanisms, progenitor and neutrino models, and particle acceleration mechanisms

Experiments

Rare metals metallicity determination, and disentangling thermal vs synchrotron emission on ~10" scales

XRISM will start probing these questions but NewAthena will give us a much sharper view.

