





XRISM science

- XRISM-Resolve: first time calorimeter are used from space
- Allows for high-spectral-resolution imaging
 - Currently at 1' level -> better with newAthena
 - Spectral resolution 5eV @ 6 keV: R~1200
 - Bummer: gate-valve not open (yet?) -> spectroscopy above ~1.8 keV
 Until now:
 - CCD spectroscopy (150 eV @ 6 keV: R~40)
 - Excellent imaging: 0.5" CXO/ACIS
 - Slitless grating spectrometers: R~1000 as well!
 - But: does not work (or difficult) for extended sources!!!
 - Resolution better at lower energies -> XRISM wins @6 keV
 - Gratings have low eff. area
- Largest impact calorimeters: extended sources!
 - supernova remnants
 - clusters of galaxies
 - diffuse ISM -> but too soft with gate-valve closed


Supernovae






- Two basic types:
 - core collapse supernovae:
 - massive stars, leave neutron star, oxygen-rich
 - thermonuclear/Type Ia supernovae:
 - % white dwarf in binary, ~0.7 Msun of iron, total disruption

Grating spectra



•LMC/SMC remnants: size small (~1') not much spectral degradation

supernova remnant shocks



- Two (or more) shocks: forward- (blastwave) and reverse snock
 - FS: shocks ISM/CSM
 - RS: shocks supernova material
- Shocks velocities: ~10,000 km/s down to ~30 km/s (t~2x10⁴ yr?)
- For ~10 30 km/s : approach sound speed ISM turbulence speeds
- Post-shock temperature (μ =0.6 average particle mass cosmic plasma):


$$\langle kT \rangle = \frac{3}{16} \mu m_{\rm p} V_{\rm s}^2 = 29 \left(\frac{\mu}{0.6}\right) \left(\frac{V_{\rm s}}{5000 \text{ km/s}}\right)^2$$


 But: electron temperatures may be lower and metal temperatures may be hotter!!

Thermal X-ray emission from SNRs

- Nonthermal emission: young SNRs with very fast shocks (V_s>3000 km/s)
- For XRISM: thermal emission more interesting:
 - Plasma optically thin (perhaps not in some lines!)
 - We see through entire SNR, front and back
 - SNRs: may expect "pure metal" plasmas (no hydrogen!)
 - Plasma is "young" and not in equilibrium:
 - non-equilibrium ionization (NEI): ionization low for kT_e
 - likely also non-equilibrium between electron, proton, metal temperatures
 - some older SNRs: reverse NEI: ionization too high for kT_e

Forward and reverse shock velocities






- Reverse shock velocity: $V_{\rm s} = \frac{dR}{dt} \frac{R}{t}$
- = shock displacement in observer frame • $\frac{dt}{dt}$ = shock displacement • $\frac{R}{R}$ free expansion velocity

Emission from hot optical thin plasma

- Radiation processes:
 - thermal bremsstrahlung $\epsilon_{\rm ff} \propto n_{\rm e} \sum_i n_{\rm i} Z_i^2 T_{\rm e}^{-1/2} \exp(-h\nu/kT_{\rm e})$
 - free-bound emission -> can dominated for pure metal plasma!
 - shape similar to bremss: but starts at edge energy!
 - two-photon continuum -> importantfor pure metal plasma!
 - line emission: depends on T_e, ionization balance, composition
- All emission depends on electron temperature, not on ion temperatures!
- All emission proportional to $\sum_i n_{\rm e} n_i V$
- \bullet standard: replace by *emission measure* $n_{\rm e}n_{\rm H}$ V (assumes that hydrogen/He dominates bremsstrahlung!)

Free-bound emission in pure metal



- In hydrogen-dominated spectrum: line emission goes up with increasing abundances
 - above certain abundances (Si ~100xsolar): also continuum goes up
 - i.e. line equivalent width stays constant above [Si]~100
- Pure metals:
 - bremsstrahlung/FB and line emission come from same species
 - free-bound becomes dominant above abundances of ~100
 - Be careful deriving densities and masses from emission measure!


NEI

• The rate change for given ion i (species),z (charge):

$$\frac{dn_{i,z}}{dt} = n_{e} \left\{ \alpha_{i,z+1}(T_{e})n_{i,z+1} + S_{i,z-1}(T_{e})n_{i,z-1} - \left[\alpha_{i,z}(T_{e}) + S_{i,z}(T_{e})n_{i,z} \right] \right\},\,$$


- $\alpha_{i,z+1}$ recombination rate from ion z+1
- $S_{i,z-1}$ ionization from z-1,
- Driven by electron collisions: $dn_{i,z}/dt \propto n_{\rm e}$
- Equilibrium: $dn_{i,z}/dt=0$ (ionizations/recombinations in balance): collisional ionization equilibrium (CIE)
- In SNRs: not enough collisions (10-26 for iron!) for equilibrium!
- The plasma is hotter, ionization lags behind!
- In some cases: overionization -> electrons have cooled but ionization has not followed!
- Relevant parameter: $\tau = n_{\rm e} t$ (ionization age, or parameter)

NEI




- ullet NEI: to relevant parameters $T_{\rm e}$ and $n_{\rm e}t$
 - More difficult to fit accurately!
 - Clusters/stars: only T_e
 - Line emission: T_e and $n_e t$; continuum: T_e (but not always reliable in case of nonthermal emission)
- Typical values: $n_{\rm e}t \approx 2 \times 10^9 \ {\rm cm}^{-3} {\rm s}$ (SN1006) to $10^{12} \ {\rm cm}^{-3} {\rm s}$ (N132D)
- In all sources (SNRs, clusters..): multiple T_e, n_et along line of sight!

He-triplets: G and R-ratios




- High resolution spectroscopy with XRISM: resolve line complexes -> He-like ions -> triplets
- Sensitive to both T_e and n_et -> break degeneracies
- Forbidden line: driven by inner shell ionization, or recombination


He-triplets: G and R-ratios



- G=(f+i)/r, R=f/i
 - issue: sometimes Li-like line contamination
- G ratio: high for severely underionized or over-ionized
- R high for underionized


Fe-K line diagnostics






- Gate-valve closed: iron K diagnostics more important
- Fe-K lines: from neutral (K α , FeI) to Fe XXV
 - Neutral Fe: 6.4 keV, fluorescence (inner shell ionization)
 - So far: centroiding
 - XRISM: getting better charge states
- In addition: getting Fe XXVI (H-like) and Ni K lines
 - With CCDs difficult to see, at edge of sensitivity
 - blending of Fe and Ni lines
 - Fe-K line ratios: evidence for nonthermal electron distributions?
 - Type Ia SNRs: Fe/Ni ratio informs us on pre-supernova conditions

What about ion temperatures?



- SNR shocks: collisionless -> may heat up electrons and ions differently!
- Extreme case: $kT_i = \frac{3}{16}m_iV_s^2$
 - i.e. iron 54xhotter than protons, electrons 1840x cooler than protons!!
- Postshock: Coulomb energy exchange
 - depends on mass ratio and charge Z
 - equilibrium scales with $n_{\rm e}t \approx 10^{12}~{\rm cm}^{-3}{\rm s}$ (same as NEI!)

What about ion temperatures?




- Ion temperature: little influence on line excitations/ionizations
- Measuring ion temperature: thermal Doppler broadening
- Equation: $kT_i = m_i \sigma_v^2$
- So measure line width -> get ion temperature
- But:
 - Also plasma velocity gradients cause lines to broaden
 - Solution: measure close to edge
 - Issue with Resolve: difficult to isolate emission from edge

What to expect from XRISM-Resolve?


- From line blends to individual lines
 - Measure triplets -> better constraints on net, kTe
 - Detect weak lines: due to resolution even weak lines stand out!
 - Higher n transitions
 - Common: abundances even elements (O,Ne,Mg, Si,S,..)
 - Lines from rare elements: Na, Al, P, Cl, K, Ti, Cr, Mn, Ni
 - some (Cr,Mn,Ni) have been detected but barely
 - Odd elements: sensitive to pre-supernova abundances, details of exposive nucleosynthesis elements
 - Currently: gate-valve will hamper detecting some of these lines


Al-K



- •Simulation for 100 ks, gatevalve open
- •Al-K line at 1.65 keV

Cas A: signature of dust grain emission





- Cas A most important young (350yr) core-collapse SNR
- Infared studies (e.g. De Looze+ 17): a lot (most?) Si contained in dust
- Nevertheless: bright Si-K lines (brightest lines!)
- How much Si in grains? And do they emit X-rays?
- Answer: yes
 - Dust sputtering: refreshing of Si ions -> low ionization Si
 - Electron/grain collisions: Si-K fluorescence lines from dust
 - Critical: dust destruction time
 - t<30 yr: most grains destroyed -> no difference
 - t>30 yr: most Si in dust -> expect signatures

Summary Or: the things I am looking forward to!

- Detecting for the first time *odd-Z elements*, and pinpointing in which ejecta layers they reside (young SNRs)
- Routine measurements of Cr, Mn, Ni abundances (Fe group)
- *Ion* temperature measurements
 - Special case: reverse shock heating; RS velocity cannot be easily measured with proper motions
- Direct evidence for pure metal plasmas: free-bound emission
- Better plasma velocity diagnostics -> can help disentangle different plasma elements along LoS
- Evidence for silicon in dust grains
- Better diagnostics of plasma elements, i.e. disentangling plasma blends